

Third Order Efficient Tests in Exponential Lattice Models

C. HIPPI

*Mathematisches Institut der Universität zu Köln,
5000 Köln 41, West Germany*

Communicated by J. Pfanzagl

In full rank multivariate exponential families of lattice distributions, one sided testing problems are considered. For these testing problems the uniformly most powerful unbiased tests are not uniformly efficient of order $o(n^{-1})$ in the class \mathcal{T}_2^* of all tests which are asymptotically similar of order $o(n^{-1})$ (Hipp, C. (1983). *J. Multivariate Anal.* **13** 67–108). In this note, better asymptotically similar tests are constructed which—for one specified value of the nuisance parameter—have maximal power in the class \mathcal{T}_2^* . The power of tests is computed up to terms of order $o(n^{-1})$, at contiguous alternatives, and for the specified value of the nuisance parameter. © 1986 Academic Press, Inc.

1. INTRODUCTION AND SUMMARY

Let p be a positive integer and v a probability measure on the Borel field \mathcal{B}^{p+1} of \mathbb{R}^{p+1} which is a lattice distribution with minimal lattice \mathbb{Z}^{p+1} . Assume that the natural parameter space

$$H^* = \left\{ \eta \in \mathbb{R}^{p+1} : \int \exp(\eta^T z) v(dz) < \infty \right\}$$

has nonvoid interior H . For $\eta \in H$ define

$$f(\eta) = \log \int \exp(\eta^T z) v(dz)$$

and P_η as the probability measure on \mathcal{B}^{p+1} with v -density

$$z \rightarrow \exp(\eta^T z - f(\eta)), \quad z \in \mathbb{R}^{p+1}.$$

Received June 18, 1984.
AMS 1980 subject classification numbers: Primary 62H15.
Key words and phrases: exponential families, lattice distributions, contiguous alternatives.

Write θ for the first component of the $(p+1)$ -vector $\eta \in H$. Fix $\theta_0 \in \mathbb{R}$ such that $T = \{\tau \in \mathbb{R}^p: (\theta_0, \tau) \in H\}$ is nonvoid.

For $n = 1, 2, \dots$ we consider the one sided testing problem

$$\{P_{\eta}^n: \eta \in H, \theta = \theta_0\} \text{ against } \{P_{\eta}^n: \eta \in H, \theta > \theta_0\}. \quad (1.1)$$

Fix a level $\alpha \in (0, 1)$. For $\tau \in T$ let $\mathcal{T}_2^*(\tau)$ denote the class of all test sequences (φ_n) with the following property: Uniformly for $u \in \mathbb{R}^p$, $|u| \leq n^{-1/2} \log n$

$$P_{(\theta_0, \tau+u)}^n \varphi_n = \alpha + o(n^{-1}) \quad (1.2)$$

and write \mathcal{T}_2^* for the class of all test sequences (φ_n) which are elements of $\mathcal{T}_2^*(\tau)$ for all $\tau \in T$.

For $\tau \in T$ write $\Sigma(\theta_0, \tau)$ for the covariance matrix of $P_{(\theta_0, \tau)}$. Partition this matrix as follows:

$$\Sigma(\theta_0, \tau) = \begin{pmatrix} \sigma_{00}(\theta_0, \tau) & \Sigma_{01}(\theta_0, \tau) \\ \Sigma_{10}(\theta_0, \tau) & \Sigma_{11}(\theta_0, \tau) \end{pmatrix}$$

where $\Sigma_{11}(\theta_0, \tau)$ is a (p, p) -matrix. Let

$$M(\theta_0, \tau) = \Sigma_{01}(\theta_0, \tau) \Sigma_{11}^{-1}(\theta_0, \tau)$$

and $T_0 = \{\tau \in T: M(\theta_0, \tau) \notin Q^p\}$.

For $n = 1, 2, \dots$ let φ_n^0 be the uniformly most powerful unbiased level α test for (1.1) (see [3, pp. 134, 135]). Then $(\varphi_n^0) \in \mathcal{T}_2^*$. It is the purpose of this note to prove the following:

THEOREM. (i) *For every $\tau_0 \in T_0$ there exists $(\varphi_n^*) \in \mathcal{T}_2^*$ with the following properties.*

(a) *For all $(\varphi_n) \in \mathcal{T}_2^*(\tau_0)$ and all $t > 0$*

$$\limsup_n n P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n (\varphi_n - \varphi_n^*) \leq 0.$$

(b) *For all $\tau \in T$ and $t > 0$*

$$\limsup_n n P_{(\theta_0 + tn^{-1/2}, \tau)}^n (\varphi_n^0 - \varphi_n^*) \leq 0.$$

(c) *For all $t > 0$*

$$\limsup_n n P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n (\varphi_n^0 - \varphi_n^*) < 0.$$

(ii) *There exists a sequence of tests (φ_n^{**}) with the following property: If $\tau_0 \in T_0$, then $(\varphi_n^{**}) \in \mathcal{T}_2^*(\tau_0)$ and for all $t > 0$*

$$\limsup_n n P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n (\varphi_n^0 - \varphi_n^{**}) < 0.$$

Remark 1. Properties (b) and (c) of (φ_n^*) yield that (φ_n^0) is—up to terms of order $o(n^{-1})$ —not asymptotically admissible in \mathcal{T}_2^* if $T_0 \neq \emptyset$.

This is true in Example 3 of [2, p. 74]. In Theorem 4 of [2, p. 81] inadmissibility of (φ_n^0) was stated under restrictive assumptions on v and α .

Remark 2. Theorem 6 of [2, p. 82] states that for all $t > 0$ and $(\varphi_n) \in \mathcal{T}_2^*$

$$P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n \varphi_n \leq H_{2n}^\alpha(t, \theta_0, \tau_0) + o(n^{-1})$$

whenever $\tau_0 \in T_0$. Here, $H_{2n}^\alpha(t, \theta_0, \tau_0)$ is the envelope power function for \mathcal{T}_2^* in the continuous case (see [4]). In our proof we shall derive the relation

$$P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n \varphi_n^* = H_{2n}^\alpha(t, \theta_0, \tau_0) + o(n^{-1}).$$

Hence $H_{2n}^\alpha(t, \theta_0, \tau_0)$ is an envelope power function for \mathcal{T}_2^* at τ_0 in the lattice case, too, provided $\tau_0 \in T_0$.

It was stated in [2, p. 72] that “randomization diminishes power.” In the construction of (φ_n^*) and (φ_n^{**}) we change the randomization of (φ_n^0) appropriately. To explain this more precisely we need some notations. Let Z_1, Z_2, \dots be independent $(p+1)$ -dimensional random vectors with distribution $P_{(\theta, \tau)}$. For $n = 1, 2, \dots$ write $Z_n = (X_n, Y_n)$ with univariate X_n and p -variate Y_n , and denote $X = X_1 + \dots + X_n$ and $Y = Y_1 + \dots + Y_n$. The tests (φ_n^0) can be written

$$\begin{aligned} \varphi_n^0 &= 1 && \text{if } X > k_n(Y, \alpha) \\ &= \gamma_n(Y, \alpha) && \text{if } X = k_n(Y, \alpha) \\ &= 0 && \text{if } X < k_n(Y, \alpha) \end{aligned}$$

with $k_n(Y, \alpha) \in \mathbb{Z}$ and $\gamma_n(Y, \alpha) \in [0, 1]$. The tests φ_n^{**} are defined as follows:

$$\begin{aligned} \varphi_n^{**} &= 1 && \text{if } X > k_n(Y, \alpha) \\ &= h(\gamma_n(Y, \alpha)) && \text{if } X = k_n(Y, \alpha) \\ &= 0 && \text{if } X < k_n(Y, \alpha). \end{aligned}$$

Here, $h: [0, 1] \rightarrow [0, 1]$ is a smooth function satisfying $h(0) = h(1) = 0$ and $\int_0^1 h(x) dx = 0$, and $h(x) < x$ if $x < \frac{1}{2}$, $h(x) > x$ if $x > \frac{1}{2}$.

The proof for $(\varphi_n^{**}) \in \mathcal{T}_2^*$ is based on the fact that $\gamma_n(Y, \alpha)$ is asymptotically uniformly distributed under $P_{(\theta_0, \tau_0)}^n$ whenever $\tau_0 \in T_0$.

The test sequence (φ_n^*) is constructed as follows: First we construct a sequence (φ_n') as above with smooth functions h_n which vary with n such that $h_n(x) \rightarrow 0$ if $x < \frac{1}{2}$, $h_n(x) \rightarrow 1$ if $x > \frac{1}{2}$.

The sequence (φ_n') is in $\mathcal{T}_2^*(\tau_0)$ and satisfies

$$P_{(\theta_0 + t n^{-1/2}, \tau_0)}^n \varphi_n' = H_{2n}^\alpha(t, \theta_0, \tau_0) + o(n^{-1}).$$

We take a \sqrt{n} -consistent estimator τ_n for τ and define φ_n^* as φ_n' if τ_n is close to τ_0 , and as φ_n^0 elsewhere.

Remark 3. If $M(\theta_0, \tau_0) \in Q^p$ then the sequence (φ_n^{**}) constructed in the proof for part (ii) of the theorem will not satisfy $(\varphi_n^{**}) \in \mathcal{T}_2^*(\tau_0)$ and

$$\limsup_n P_{(\theta_0 + t n^{-1/2}, \tau_0)}^n (\varphi_n^0 - \varphi_n^{**}) < 0 \quad \text{for all } t > 0$$

in general. In Example 2 of [2, p. 73] the sequence (φ_n^0) satisfies

$$\liminf_n P_{(\theta_0 + t n^{-1/2}, \tau_0)}^n (\varphi_n^0 - \varphi_n) \geq 0$$

for all $t > 0$ and $(\varphi_n) \in \mathcal{T}_2^*(\tau_0)$ if $\alpha = \frac{1}{2}$, and

$$\limsup_n P_{(\theta_0 + t n^{-1/2}, \tau_0)}^n (\varphi_n^0 - \varphi_n) \geq 0$$

for all $t > 0$ and $(\varphi_n) \in \mathcal{T}_2^*(\tau_0)$ for arbitrary α .

Remark 4. Let $M(\theta_0, \tau_0) = (1/m)(k_1, \dots, k_p)$ with relatively prime integers m, k_1, \dots, k_p . Then the distribution of $\gamma_n(Y, \alpha)$ can be approximated by $\delta(a_n) * U * V$, where $a_n \in \mathbb{R}$, $\delta(a_n)$ is the Dirac measure at a_n , U is the uniform distribution on $\{0, 1/m, \dots, (m-1)/m\}$, and V is a noncentral chi-square distribution; the convolution is computed on the torus \mathbb{R}/\mathbb{Z} . The constants a_n oscillate, and the oscillation depends on the value of the nuisance parameter τ_0 . A function h yielding a sequence (φ_n^{**}) satisfying $(\varphi_n^{**}) \in \mathcal{T}_2^*(\tau_0)$ has to satisfy the conditions $Eh^*(U+x) = EUh^{**}(U+x) = 0$ for all x , where $h^*(x) = h(x) - x$. These conditions, however, imply $h^* \equiv 0$.

2. PROOF OF THE THEOREM

In this section we shall use some notations of [2] without repeating the definitions.

For $j = 1, 2, \dots$ let $h_j^*: [0, 1] \rightarrow [0, 1]$ be a trigonometric polynomial of the form

$$h_j^*(x) = \sum \{a_l \exp(2\pi i l x) : 0 < |l| \leq j\}$$

with $h_j^*(0) = 0$.

Assume that $h_j^*(x) < 0$ for $0 < x < \frac{1}{2}$, $h_j^*(x) > 0$ for $\frac{1}{2} < x < 1$, and that for all $0 < \varepsilon < \frac{1}{2}$, $h_j^*(x) \rightarrow -x$, uniformly for $\varepsilon < x < \frac{1}{2} - \varepsilon$, and $h_j^*(x) \rightarrow 1 - x$, uniformly for $\frac{1}{2} + \varepsilon < x < 1 - \varepsilon$. Let $h_j(x) = h_j^*(x) + x$. For $n = 2, 3, \dots$ let $\tau_n(Z_1, \dots, Z_n)$ be an estimator for τ such that uniformly for (θ, τ) in compact subsets of H

$$P_{(\theta, \tau)}^n \{ |\tau_n(Z_1, \dots, Z_n) - \tau| \geq n^{-1/2} \log n \} = o(n^{-1}).$$

To simplify our notation we shall omit arguments whenever possible. Our first result yields a stochastic expansion for $\gamma_n(Y, \alpha)$.

LEMMA 1. *We have uniformly for τ in compact subsets of T and $y \in A(n, (\theta_0, \tau))$*

$$\begin{aligned} \gamma_n(y, \alpha) = & c + Q_1(v, \tilde{y}(n, \theta_0, \tau), \theta_0, \tau) \\ & + n^{-1/2} \{ -Q_2(v, \tilde{y}(n, \theta_0, \tau), \theta_0, \tau) \\ & + Q_1(v, \tilde{y}(n, \theta_0, \tau), \theta_0, \tau) W_1(v, \tilde{y}(n, \theta_0, \tau), \theta_0, \tau) \\ & + \frac{1}{2} \sigma^{-1} N_x \gamma_n^2(y, \alpha) \\ & - \sigma^{-1} N_x \gamma_n(y, \alpha) Q_1(v, \tilde{y}(n, \theta_0, \tau), \theta_0, \tau) \} + o(n^{-1/2}). \end{aligned}$$

Recall that

$$c = c(y) = k_n(y, \alpha) - n\mu_0 + nM\mu_1 + n^{1/2}\sigma N_x - My$$

and

$$v = v(y) = -\sigma N_x + M\tilde{y}(n, \theta_0, \tau),$$

and that $Q_1(v, \tilde{y}(n, \theta_0, \tau), \theta_0, \tau)$ is a polynomial in $\tilde{y}(n, \theta_0, \tau)$ of degree two or less.

LEMMA 2. *For $\tau \in T$ let $P(y, \tau)$ be a polynomial in $y \in \mathbb{R}^p$ such that the degree and the coefficients of P remain bounded as long as τ varies in compact subsets of T . Then for all $\tau \in T_0$ and $m = 1, 2, \dots$ we have uniformly for $|u| \leq n^{-1/2} \log n$*

$$\begin{aligned} P_{(\theta_0, \tau+u)}^n [& h_m^*(c(Y) + Q_1(v(Y), \tilde{Y}(n, \theta_0, \tau+u), \theta_0, \tau+u)) \\ & \times P(\tilde{Y}(n, \theta_0, \tau+u), \tau+u)] = o(n^{-1/2}). \end{aligned}$$

Proof. It suffices to prove the assertion for

$$h_m^*(x) = \exp(2\pi i m x), \quad x \in \mathbb{R}, m \neq 0.$$

For $w \in \mathbb{R}^p$ consider

$$f(w) = P_{(\theta_0, \tau + u)}^n h_m^*(c(Y) + w^T \tilde{Y}(n, \theta_0, \tau + u)).$$

With

$$\begin{aligned} H(n) &= n^{1/2} \sigma(\theta_0, \tau + u) N_\alpha - n \mu_0(\theta_0, \tau + u) \\ &\quad + n M(\theta_0, \tau + u) \mu_1(\theta_0, \tau + u) \end{aligned}$$

we obtain

$$\begin{aligned} f(w) &= P_{(\theta_0, \tau + u)}^n h_m^*((-M(\theta_0, \tau + u) + n^{-1/2} w)^T Y \\ &\quad + H(n) - n^{1/2} w^T \mu_1(\theta_0, \tau + u)) \\ &= (P_{(\theta_0, \tau + u)} h_m^*((-M(\theta_0, \tau + u) + n^{-1/2} w)^T Y_1))^n \\ &\quad \times h_m^*(H(n) - n^{1/2} w^T \mu_1(\theta_0, \tau + u)). \end{aligned}$$

There exists a positive ε such that for $|u| \leq \varepsilon$ and $|W| \leq \varepsilon$ we have

$$|P_{(\theta_0, \tau + u)} h_m^*((-M(\theta_0, \tau + u) + W)^T Y_1)| \leq 1 - \varepsilon.$$

This implies that for all $s > 0$ and uniformly for $|u| \leq n^{-1/2} \log n$ and $|w| \leq \varepsilon n^{1/2}$

$$f(w) = o(n^{-s}). \quad (2.1)$$

The same is true for derivatives of all orders:

If β_1, \dots, β_p are nonnegative integers, then for all $s > 0$ and uniformly for $|u| \leq n^{-1/2} \log n$ and $|w| \leq \varepsilon n^{1/2}$

$$\frac{\partial^{\beta_1 + \dots + \beta_p}}{\partial w_1^{\beta_1} \dots \partial w_p^{\beta_p}} f(w) = o(n^{-s}).$$

Let $K|_{\mathcal{B}^p}$ be a probability measure satisfying $\int |x|^2 K(dx) < \infty$ which has a characteristic function $\tilde{K}(t)$ vanishing for $|t| > 1$. If g is a measurable function which is bounded by some polynomial, then (2.1) and (2.2) together with Lemma 11.6 in [1, p. 98] imply that for all $\tau \in T$ and $s > 0$ and uniformly for $|u| \leq n^{-1/2} \log n$

$$\int P_{(\theta_0, \tau + u)}^n h_m^*(c(Y)) g(\tilde{Y}(n, \theta_0, \tau + u) + \varepsilon^{-1} n^{-1/2} w) K(dw) = o(n^{-s}).$$

Let $G: \mathbb{R}^p \rightarrow \mathbb{R}$ be the polynomial of degree two or less such that $G(\tilde{Y}(n, \theta_0, \tau + u)) = Q_1(v(y), \tilde{Y}(n, \theta_0, \tau + u), \theta_0, \tau + u)$, and let $g_m(y) = h_m^*(G(y)) P(y, \tau + u)$. Then for all nonnegative integers β_1, \dots, β_p there exists a polynomial Q in y such that for all $u \in \mathbb{R}^p$ with $|u| \leq n^{-1/2} \log n$

$$\left| \frac{\partial^{\beta_1 + \dots + \beta_p}}{\partial y_1^{\beta_1} \dots \partial y_p^{\beta_p}} g_m(y) \right| \leq k^{\beta_1 + \dots + \beta_p} Q(y).$$

This implies that for all $s > 0$ and nonnegative integers β_1, \dots, β_p we have uniformly for $|u| \leq n^{-1/2} \log n$

$$\begin{aligned} & \int P_{(\theta_0, \tau + u)}^n h_m^*(c(Y)) \frac{\partial^{\beta_1 + \dots + \beta_p}}{\partial y_1^{\beta_1} \dots \partial y_p^{\beta_p}} g_m(y) \Big|_{Y = \tilde{Y}(n, \theta_0, \tau + u) + \varepsilon^{-1} n^{-1/2} w} K(dw) \\ &= o(n^{-s}). \end{aligned}$$

The remainder $R_n(x, w)$ in the Taylor expansion

$$\begin{aligned} g_m(x) &= \sum_{0 \leq \beta_1 + \dots + \beta_p \leq 1} (-\varepsilon n^{1/2})^{-(\beta_1 + \dots + \beta_p)} w_1^{\beta_1} \dots w_p^{\beta_p} \\ &\quad \times \frac{\partial^{\beta_1 + \dots + \beta_p}}{\partial y_1^{\beta_1} \dots \partial y_p^{\beta_p}} g_m(y) \Big|_{Y = x + \varepsilon^{-1} n^{-1/2} w} \prod_{j=1}^p (\beta_j!)^{-1} + R_n(x, w) \end{aligned}$$

satisfies

$$R_n(x, w) \leq n^{-1} |w|^2 R(x),$$

where R is a polynomial which may depend on τ but not on u . Integrating both sides of the Taylor expansion with respect to $P_{(\theta_0, \tau + u)}^n$ and K we obtain that uniformly for $|u| \leq n^{-1/2} \log n$

$$P_{(\theta_0, \tau + u)}^n h_m^*(c(Y)) g_m(\tilde{Y}(n, \theta_0, \tau + u)) = o(n^{-1/2}).$$

This implies the assertion of Lemma 2.

For fixed m define the sequence of tests (φ_n) by

$$\begin{aligned} \varphi_n &= 1 & \text{if } X > k_n(Y, \alpha) \\ &= h_m(\gamma_n(Y, \alpha)) & \text{if } X = k_n(Y, \alpha) \\ &= 0 & \text{if } X < k_n(Y, \alpha). \end{aligned}$$

LEMMA 3. For $\tau_0 \in T_0$ we have $(\varphi_n) \in \mathcal{T}_2^*(\tau_0)$.

Proof. We have to show that uniformly for $|u| \leq n^{-1/2} \log n$

$$P_{(\theta_0, \tau_0 + u)}^n h_m^*(\gamma_n(Y, \alpha)) 1_{\{X = k_n(Y, \alpha)\}} = o(n^{-1}).$$

Recall that uniformly for $|u| \leq n^{-1/2} \log n$

$$P_{(\theta_0, \tau_0 + u)}^n \{Y \notin A(n, (\theta_0, \tau_0 + u))\} = o(n^{-1}).$$

With Lemma 1 we obtain that uniformly for $|u| \leq n^{-1/2} \log n$

$$\begin{aligned} & P_{(\theta_0, \tau_0 + u)}^n h_m^*(\gamma_n(Y, \alpha)) 1_{\{X = k_n(Y, \alpha)\}} \\ &= n^{-1/2} \sigma^{-1}(\theta_0, \tau_0 + u) \varphi(N_\alpha) P_{(\theta_0, \tau_0 + u)}^n \left[\left\{ 1 + n^{-1/2} \sigma^{-1} N_\alpha \gamma_n(Y, \alpha) \right. \right. \\ &\quad \left. \left. - n^{-1/2} \sigma^{-1} N_\alpha Q_1(v(Y), \tilde{Y}(n, \theta_0, \tau_0 + u), \tau_0) \right. \right. \\ &\quad \left. \left. + n^{-1/2} W_1(v(Y), \tilde{Y}(n, \theta_0, \tau_0 + u), \theta_0, \tau_0) \right\} \right. \\ &\quad \left. \times \{h_m^*(c(Y) + Q_1(v(Y), \tilde{Y}(n, \theta_0, \tau_0 + u), \tau_0 + u)) \right. \\ &\quad \left. + h_m^*(c(Y) + Q_1(v(Y), \tilde{Y}(n, \theta_0, \tau_0 + u), \tau_0)) \right. \\ &\quad \left. \times n^{-1/2} [-Q_2(v(Y), \tilde{Y}(n, \theta_0, \tau_0 + u), \theta_0, \tau_0) \right. \\ &\quad \left. + Q_1(v(Y), \tilde{Y}(n, \theta_0, \tau_0 + u), \tau_0) W_1(v(Y), \tilde{Y}(n, \theta_0, \tau_0 + u), \theta_0, \tau_0) \right. \\ &\quad \left. + \frac{1}{2} \sigma^{-1}(\theta_0, \tau_0) N_\alpha \gamma_n^2(Y, \alpha) \right. \\ &\quad \left. \left. - \sigma^{-1}(\theta_0, \tau_0) N_\alpha \gamma_n(Y, \alpha) Q_1(v(Y), \tilde{Y}(n, \theta_0, \tau_0 + u), \tau_0) \right\} \right] \\ &\quad + o(n^{-1}). \end{aligned}$$

From Lemma 2 we obtain that uniformly for $|u| \leq n^{-1/2} \log n$

$$\begin{aligned} & P_{(\theta_0, \tau_0 + u)}^n h_m^*(\gamma_n(Y, \alpha)) 1_{\{X = k_n(Y, \alpha)\}} \\ &= n^{-1} \sigma^{-1}(\theta_0, \tau_0 + u) \varphi(N_\alpha) P_{(\theta_0, \tau_0 + u)}^n \left[\sigma^{-1} N_\alpha \gamma_n(Y, \alpha) h_m^*(\gamma_n(Y, \alpha)) \right. \\ &\quad \left. + \frac{1}{2} \sigma^{-1} N_\alpha \gamma_n^2(Y, \alpha) h_m^{*'}(\gamma_n(Y, \alpha)) \right. \\ &\quad \left. - \sigma^{-1} N_\alpha \gamma_n(Y, \alpha) Q_1(v(Y), \tilde{Y}(n, \theta_0, \tau_0 + u), \tau_0) h_m^{*'}(\gamma_n(Y, \alpha)) \right] \\ &\quad + o(n^{-1}). \end{aligned}$$

In the proof of Lemma 2 we have seen that for nonzero integers k and for $w \in \mathbb{R}^p$ and uniformly for $|u| \leq n^{-1/2} \log n$

$$P_{(\theta_0, \tau_0 + u)}^n \exp(2\pi i k \gamma_n(Y, \alpha) + i w^T \tilde{Y}(n, \theta_0, \tau_0 + u)) = o(n^0).$$

This implies that for $F: [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ bounded, continuous, $F(0, x) = F(1, x)$ for all $x \in \mathbb{R}$, we have uniformly for $|u| \leq n^{-1/2} \log n$

$$P_{(\theta_0, \tau_0 + u)}^n F(\gamma_n(Y, \alpha), \tilde{Y}(n, \theta_0, \tau_0 + u)) = \int_0^1 F(x, y) \varphi_{\Sigma_{11}}(y) dy dx + o(n^0).$$

Hence uniformly for $|u| \leq n^{-1/2} \log n$

$$\begin{aligned} P_{(\theta_0, \tau_0 + u)}^n h_m^*(\gamma_n(Y, \alpha)) 1_{\{X = k_n(Y, \alpha)\}} \\ = n^{-1} \sigma^{-1}(\theta_0, \tau_0 + u) \varphi(N_\alpha) \left\{ \sigma^{-1} N_\alpha \int_0^1 (x h_m^*(x) + \frac{1}{2} x^2 h_m^{*'}(x)) dx \right. \\ \left. - \sigma^{-1} N_\alpha \int_0^1 x h_m^{*'}(x) G(y) \varphi_{\Sigma_{11}}(y) dy dx \right\} + o(n^{-1}). \end{aligned}$$

Here, G is the polynomial of degree two or less satisfying

$$G(\tilde{y}(n, \theta_0, \tau_0 + u)) = Q_1(v(y), \tilde{y}(n, \theta_0, \tau_0 + u), \tau_0).$$

The term involving G has an unbounded smooth integrand. Here, a simple truncation argument is needed.

The assertion of Lemma 3 now follows from

$$\int_0^1 x h_m^{*'}(x) dx = 0$$

and

$$\int_0^1 (x h_m^*(x) + \frac{1}{2} x^2 h_m^{*'}(x)) dx = 0.$$

Our next result is a modification of the Corollary in [2, p. 94]. Its proof is essentially the same as the one given in [2].

LEMMA 4. For $y \in \mathbb{Z}^p$ let $\alpha_n(y) \in (0, 1)$ be given and define $\gamma_n(y) \in [0, 1)$ and $k_n(y) \in \mathbb{Z}$ by

$$Q(n, y, \theta_0) \{w > k_n(y)\} + \gamma_n(y) Q(n, y, \theta_0) \{k_n(y)\} = \alpha_n(y),$$

and define

$$\begin{aligned} \varphi_n &= 1 && \text{if } X > k_n(Y) \\ &= \gamma_n(Y) && \text{if } X = k_n(Y) \\ &= 0 && \text{if } X < k_n(Y). \end{aligned}$$

Fix $\tau_0 \in T$ and assume that uniformly for τ with $|\tau - \tau_0| \leq n^{-1/2} \log n$ and $y \in A(n, (\theta_0, \tau))$

$$\alpha_n(y) - \alpha = o(n^{-1/3}).$$

If $\varphi_n \in \mathcal{T}_2^*(\tau_0)$, then uniformly for t in compact subsets of \mathbb{R}

$$P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n(\varphi_n - \varphi_n^0) = I_2 + o(n^{-1})$$

where I_2 is defined in Lemma 2 of [2, p. 92].

Remark 5. Theorem 6 in [2, p. 82] states that for $\tau_0 \in T_0$ we have for all $(\varphi_n) \in \mathcal{T}_2^*$ and $t > 0$

$$P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n \varphi_n \leq H_{2n}^a(t, \theta_0, \tau_0) + o(n^{-1}).$$

From the proof given there we see that this inequality holds for all $(\varphi_n) \in \mathcal{T}_2^*(\tau_0)$.

To prove part (ii) of the theorem, fix $m \geq 1$ and define

$$\begin{aligned} \varphi_n^{**} &= 1 && \text{if } X > k_n(Y, \alpha) \\ &= h_m(\gamma_n(Y, \alpha)) && \text{if } X = k_n(Y, \alpha) \\ &= 0 && \text{if } X < k_n(Y, \alpha). \end{aligned}$$

Lemma 3 yields $(\varphi_n^{**}) \in \mathcal{T}_2^*(\tau_0)$ for all $\tau_0 \in T_0$. Lemma 4 implies that for all $\tau_0 \in T$ with $M(\theta_0, \tau_0) \notin Q^p$ and $t > 0$

$$P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n(\varphi_n^{**} - \varphi_n^0) = n^{-1}t\sigma^{-1}\varphi(N_x + t\sigma) \int_0^1 h_m^*(x)(x - \tfrac{1}{2}) dx + o(n^{-1}).$$

The integral is positive since $h_m^*(x) < 0$ for $x < \frac{1}{2}$, $h_m^*(x) > 0$ for $x > \frac{1}{2}$.

To prove part (i) consider the family of test sequences $(\varphi_n^{(m)}) \in \mathcal{T}_2^*(\tau_0)$ constructed with h_m . Let $m = m(n)$ tend to infinity with n such that the sequence (φ_n') , $\varphi_n' = \varphi_n^{(m(n))}$, is still an element of $\mathcal{T}_2^*(\tau_0)$.

Let $\varphi_n^* = \varphi_n'$ if $|\tau_n - \tau_0| \leq 2n^{-1/2} \log n$, and $\varphi_n^* = \varphi_n^0$ elsewhere. Then $(\varphi_n^*) \in \mathcal{T}_2^*$. To see this let $\tau \in T$ be arbitrary. If $\tau \neq \tau_0$, then uniformly for $|u| \leq n^{1/2} \log n$

$$P_{(\theta_0, \tau + u)}^n \{|\tau_n - \tau_0| \leq 2n^{-1/2} \log n\} = o(n^{-1})$$

and therefore

$$P_{(\theta_0, \tau + u)}^n \varphi_n^* = P_{(\theta_0, \tau + u)}^n \varphi_n^0 + o(n^{-1}) = \alpha + o(n^{-1}).$$

If $\tau = \tau_0$, then uniformly for $|u| \leq n^{-1/2} \log n$

$$P_{(\theta_0, \tau_0 + u)}^n \{|\tau_n - \tau_0 - u| > n^{-1/2} \log n\} = o(n^{-1})$$

and therefore

$$P_{(\theta_0, \tau_0 + u)}^n \{|\tau_n - \tau_0| > 2n^{-1/2} \log n\} = o(n^{-1}).$$

This yields that uniformly for $|u| \leq n^{-1/2} \log n$

$$P_{(\theta_0, \tau_0 + u)}^n \varphi_n^* = P_{(\theta_0, \tau_0 + u)}^n \varphi_n' + o(n^{-1}) = \alpha + o(n^{-1}).$$

For $\tau \in T$ with $\tau \neq \tau_0$ we obtain from

$$P_{(\theta_0 + tn^{-1/2}, \tau)}^n \{|\tau_n - \tau| > n^{-1/2} \log n\} = o(n^{-1})$$

that

$$P_{(\theta_0 + tn^{-1/2}, \tau)}^n (\varphi_n^* - \varphi_n^0) = o(n^{-1}).$$

Furthermore, for $t > 0$ we have

$$\begin{aligned} P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n \varphi_n^* &= P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n \varphi_n' + o(n^{-1}) \\ &= P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n \varphi_n^0 + n^{-1} t \sigma^{-1} \varphi(N_\alpha + t\sigma) \\ &\quad \times P_{(\theta_0 + tn^{-1/2}, \tau)}^n h_{m(n)}^*(\gamma_n(Y, \alpha))(\gamma_n(Y, \alpha) - \tfrac{1}{2}) + o(n^{-1}). \end{aligned}$$

For $0 < \varepsilon < \frac{1}{2}$ let $B(n, \alpha)$ be the set of all y for which $\gamma_n(y, \alpha)$ is an element of $[0, \varepsilon)$ or $(1 - \varepsilon, 1)$ or $(\frac{1}{2} - \varepsilon, \frac{1}{2} + \varepsilon)$. Then for n sufficiently large, $h_{m(n)}(\gamma_n(y, \alpha)) < \varepsilon$ if $y \notin B(n, \varepsilon)$ and $\gamma_n(y, \alpha) < \frac{1}{2}$, and $h_{m(n)}(\gamma_n(y, \alpha)) > 1 - \varepsilon$ if $y \notin B(n, \varepsilon)$ and $\gamma_n(y, \alpha) > \frac{1}{2}$. Hence, with

$$\begin{aligned} h_0^*(x) &= -x, & x \leq \tfrac{1}{2} \\ &= 1 - x, & x > \tfrac{1}{2} \end{aligned}$$

we have for n sufficiently large

$$|h_{m(n)}^*(\gamma_n(y, \alpha)) - h_0^*(\gamma_n(y, \alpha))| \leq \varepsilon + 21_{B(n, \varepsilon)}.$$

This implies that for n sufficiently large

$$\begin{aligned} &|P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n h_{m(n)}^*(\gamma_n(Y, \alpha))(\gamma_n(Y, \alpha) - \tfrac{1}{2}) \\ &\quad - P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n h_0^*(\gamma_n(Y, \alpha))(\gamma_n(Y, \alpha) - \tfrac{1}{2})| \\ &\leq \varepsilon + 2P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n \{Y \in B(n, \varepsilon)\}. \end{aligned}$$

Since under $P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n$, $\gamma_n(Y, \alpha)$ is asymptotically uniformly distributed on $[0, 1]$, we have

$$\lim_n P_{(\theta_0 + tn^{-1/2}, \tau)}^n \{Y \in B(n, \varepsilon)\} = 4\varepsilon$$

and

$$\lim_n P_{(\theta_0 + tn^{-1/2}, \tau)}^n h_0^*(\gamma_n(Y, \alpha))(\gamma_n(Y, \alpha) - \tfrac{1}{2}) = \tfrac{1}{24}.$$

Since $\varepsilon > 0$ can be chosen arbitrarily small, we obtain

$$\lim_n P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n h_{m(n)}^*(\gamma_n(Y, \alpha))(\gamma_n(Y, \alpha) - \frac{1}{2}) = \frac{1}{24}$$

or

$$\lim_n P_{(\theta_0 + tn^{-1/2}, \tau_0)}^n \varphi_n^* = H_{2n}^\alpha(t, \theta_0, \tau_0) + o(n^{-1}).$$

According to our Remark 5, $H_{2n}^\alpha(t, \theta_0, \tau_0)$ is—up to errors of order $o(n^{-1})$ —an upper bound for the power of tests in $\mathcal{T}_2^*(\tau_0)$. This proves property (b) of (φ_n^*) . Property (c) of (φ_n^*) now follows from Theorem 2(iv) of [2, p. 78].

REFERENCES

- [1] BHATTACHARYA, R. N., AND RANGA RAO, R. (1976). *Normal Approximations and Asymptotic Expansions*. Wiley, New York.
- [2] HIPPI, C. (1983). Third order efficiency of conditional tests in exponential models: the lattice case. *J. Multivariate Anal.* **13** 67–108.
- [3] LEHMANN, E. L. (1959). *Testing Statistical Hypotheses*. Wiley, New York.
- [4] MICHEL, R. (1979). Third order efficiency of conditional tests for exponential families. *J. Multivariate Anal.* **9** 401–409.